

The American Psychiatric Association Practice Guideline for the Treatment of Patients With Eating Disorders

Catherine Crone, M.D. (Chair), Laura J. Fochtmann, M.D., M.B.I. (Vice-Chair, Methodologist), Evelyn Attia, M.D., Robert Boland, M.D., Javier Escobar, M.D., Victor Fornari, M.D., Neville Golden, M.D., Angela Guarda, M.D., Maga Jackson-Triche, M.D., Laurie Manzo, M.Ed., R.D., L.D.N., Margherita Mascolo, M.D., Karen Pierce, M.D., Megan Riddle, M.D., Ph.D., M.S., Andreea Seritan, M.D., Blair Uniacke, M.D., Nancy Zucker, Ph.D., Systematic Review: Joel Yager, M.D., Thomas J. Craig, M.D., Seung-Hee Hong, Jennifer Medicus

At its April 2021 meeting, the American Psychiatric Association (APA) Board of Trustees approved “The American Psychiatric Association Practice Guideline for the Treatment of Patients With Eating Disorders.” The full guideline is available at APA’s Practice Guidelines website.

The goal of this guideline is to improve the quality of care and treatment outcomes for patients with eating disorders, as defined by the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Text Revision (DSM-5-TR; American Psychiatric Association 2022). Since publication of the last American Psychiatric Association (APA) practice guideline on eating disorders (American Psychiatric Association 2006), there have been many studies on psychotherapies for individuals with these diagnoses as well as some studies on pharmacotherapies. Despite this, there are still substantial gaps in the availability and use of evidence-based treatments for individuals with an eating disorder (Kazdin et al. 2017). This practice guideline aims to help clinicians enhance care for their patients by reviewing current evidence and providing evidence-based statements (Box 1) that are intended to increase knowledge, improve assessment, and optimize treatment of eating disorders.

The lifetime prevalence of eating disorders in the United States is approximately 0.80% for anorexia nervosa (AN), 0.28% for bulimia nervosa (BN), and 0.85% for binge-eating disorder (BED) (Udo and Grilo 2018), although estimates can vary depending on the study location, sample demographic characteristics, case finding, and diagnostic approaches (Galmiche et al. 2019; Santomauro et al. 2021; Wu et al. 2020). Furthermore, data suggest an increasing incidence of eating disorders and inpatient care for eating disorders, particularly AN, during the COVID-19 pandemic (Agostino et al. 2021; Asch et al. 2021; Otto et al. 2021; Taquet et al. 2021). Importantly, the lifetime burdens and psychosocial impairments associated with an eating disorder can be substantial because these illnesses can persist for decades, and they typically have an onset in adolescence or early adulthood (Udo and Grilo 2018).

In the United States, for the 2018–2019 fiscal year, the total economic costs of eating disorders were estimated to be \$64.7 billion, with an additional \$326.5 billion

attributable to reductions in well-being associated with eating disorders (Streatfeild et al. 2021).

Eating disorders are also associated with increases in all-cause mortality and deaths due to suicide (Auger et al. 2021; Nielsen and Vilmar 2021; Tith et al. 2020; van Hoeken and Hoek 2020). In addition, rates of suicide attempts are increased in individuals who have an eating disorder (Keski-Rahkonen 2021; Smith et al. 2018; Udo et al. 2019). Morbidity and mortality among individuals with an eating disorder are heightened by the common co-occurrence of health conditions such as diabetes and other psychiatric disorders, particularly depression, anxiety, posttraumatic stress disorder (PTSD), obsessive-compulsive disorder (OCD), ADHD, and substance use disorders (Ahn et al. 2019; Cliffe et al. 2020; Gibbings et al. 2021; Keski-Rahkonen 2021; Udo and Grilo 2019).

Accordingly, the overall goal of this guideline is to enhance the assessment and treatment of eating disorders, thereby reducing the mortality, morbidity, and significant psychosocial and health consequences of these important psychiatric conditions.

OVERVIEW OF THE DEVELOPMENT PROCESS

Since the publication of Clinical Practice Guidelines We Can Trust (Institute of Medicine 2011), a report of the Institute of Medicine (now known as National Academy of Medicine), there has been an increasing focus on using clearly defined, transparent processes for rating the quality of evidence and the strength of the overall body of evidence in systematic reviews of the scientific literature. This guideline was developed using a process intended to be consistent with the recommendations of the Institute of Medicine (Institute of Medicine 2011) and the Principles for the Development of Specialty Society Clinical Guidelines of the Council of Medical Specialty Societies (2012). Parameters used for the guideline’s systematic review are included with the full text of the guideline. The APA website features a full description of the guideline development process.

Box 1. Guideline Statements^a

Assessment and Determination of Treatment Plan

1. APA recommends **(1C)** screening for the presence of an eating disorder as part of an initial psychiatric evaluation.
2. APA recommends **(1C)** that the initial evaluation of a patient with a possible eating disorder include assessment of
 - the patient's height and weight history (e.g., maximum and minimum weight, recent weight changes);
 - presence of, patterns in, and changes in restrictive eating, food avoidance, binge eating, and other eating-related behaviors (e.g., rumination, regurgitation, chewing and spitting);
 - patterns and changes in food repertoire (e.g., breadth of food variety, narrowing or elimination of food groups);
 - presence of, patterns in, and changes in compensatory and other weight control behaviors, including dietary restriction, compulsive or driven exercise, purging behaviors (e.g., laxative use, self-induced vomiting), and use of medication to manipulate weight;
 - percentage of time preoccupied with food, weight, and body shape;
 - prior treatment and response to treatment for an eating disorder;
 - psychosocial impairment secondary to eating or body image concerns or behaviors; and
 - family history of eating disorders, other psychiatric illnesses, and other medical conditions (e.g., obesity, inflammatory bowel disease, diabetes mellitus).
3. APA recommends **(1C)** that the initial psychiatric evaluation of a patient with a possible eating disorder include weighing the patient and quantifying eating and weight control behaviors (e.g., frequency, intensity, or time spent on dietary restriction, binge eating, purging, exercise, and other compensatory behaviors).
4. APA recommends **(1C)** that the initial psychiatric evaluation of a patient with a possible eating disorder identify co-occurring health conditions, including co-occurring psychiatric disorders.
5. APA recommends **(1C)** that the initial psychiatric evaluation of a patient with a possible eating disorder include a comprehensive review of systems.
6. APA recommends **(1C)** that the initial physical examination of a patient with a possible eating disorder include assessment of vital signs, including temperature, resting heart rate, blood pressure, orthostatic pulse, and orthostatic blood pressure; height, weight, and BMI (or percent median BMI, BMI percentile, or BMI Z-score for children and adolescents); and physical appearance, including signs of malnutrition or purging behaviors.
7. APA recommends **(1C)** that the laboratory assessment of a patient with a possible eating disorder include a complete blood count and a comprehensive metabolic panel, including electrolytes, liver enzymes, and renal function tests.
8. APA recommends **(1C)** that an electrocardiogram be done in patients with a restrictive eating disorder, patients with severe purging behavior, and patients who are taking medications that are known to prolong QTc intervals.
9. APA recommends **(1C)** that patients with an eating disorder have a documented, comprehensive, culturally appropriate,

and person-centered treatment plan that incorporates medical, psychiatric, psychological, and nutritional expertise, commonly via a coordinated multidisciplinary team.

Anorexia Nervosa

10. APA recommends **(1C)** that patients with anorexia nervosa who require nutritional rehabilitation and weight restoration have individualized goals set for weekly weight gain and target weight.
11. APA recommends **(1B)** that adults with anorexia nervosa be treated with an eating disorder-focused psychotherapy, which should include normalizing eating and weight control behaviors, restoring weight, and addressing psychological aspects of the disorder (e.g., fear of weight gain, body image disturbance).
12. APA recommends **(1B)** that adolescents and emerging adults with anorexia nervosa who have an involved caregiver be treated with eating disorder-focused family based treatment, which should include caregiver education aimed at normalizing eating and weight control behaviors and restoring weight.

Bulimia Nervosa

13. APA recommends **(1C)** that adults with bulimia nervosa be treated with eating disorder-focused cognitive-behavioral therapy and that a serotonin reuptake inhibitor (e.g., 60 mg fluoxetine daily) also be prescribed, either initially or if there is minimal or no response to psychotherapy alone by 6 weeks of treatment.
14. APA suggests **(2C)** that adolescents and emerging adults with bulimia nervosa who have an involved caregiver be treated with eating disorder-focused family based treatment.

Binge-Eating Disorder

15. APA recommends **(1C)** that patients with binge-eating disorder be treated with eating disorder-focused cognitive-behavioral therapy or interpersonal therapy, in either individual or group formats.
16. APA suggests **(2C)** that adults with binge-eating disorder who prefer medication or have not responded to psychotherapy alone be treated with either an antidepressant medication or lisdexamfetamine.

^aThe authors of the guideline determined each final rating, as described in the section "Guideline Development Process" (see Table 1 in the full guideline). A recommendation (denoted by the numeral 1 after the guideline statement) indicates confidence that the benefits of the intervention clearly outweigh harms. A suggestion (denoted by the numeral 2 after the guideline statement) indicates greater uncertainty. Although the benefits of the statement are still viewed as outweighing the harms, the balance of benefits and harms is more difficult to judge, or either the benefits or the harms may be less clear. With a suggestion, patient values and preferences may be more variable, and this can influence the clinical decision that is ultimately made. Each guideline statement also has an associated rating for the strength of supporting research evidence. Three ratings are used: high, moderate, and low (denoted by the letters A, B, and C, respectively) and reflect the level of confidence that the evidence for a guideline statement reflects a true effect based on consistency of findings across studies, directness of the effect on a specific health outcome, precision of the estimate of effect, and risk of bias in available studies (Agency for Healthcare Research and Quality 2014; Balshem et al. 2011; Guyatt et al. 2006).

RATING THE STRENGTH OF RESEARCH EVIDENCE AND RECOMMENDATIONS

Development of guideline statements entails weighing the potential benefits and harms of the statement and then identifying the level of confidence in that determination. (See Appendix G in the supplemental information accompanying the full guideline online for detailed descriptions of the potential benefits and harms for each statement.) This concept of balancing benefits and harms to determine guideline recommendations and strength of recommendations is a hallmark of GRADE (Grading of Recommendations Assessment, Development and Evaluation), which is used by multiple professional organizations around the world to develop practice guideline recommendations (Guyatt et al. 2013). With the GRADE approach, recommendations are rated by assessing the confidence that the benefits of the statement outweigh the harms and burdens of the statement, determining the confidence in estimates of effect as reflected by the quality of evidence, estimating patient values and preferences (including whether they are similar across the patient population), and identifying whether resource expenditures are worth the expected net benefit of following the recommendation (Andrews et al. 2013).

In weighing the balance of benefits and harms for each statement in this guideline, our level of confidence is informed by available evidence (see Appendix C in the supplemental information accompanying the full guideline online), which includes evidence from clinical trials as well as expert opinion and patient values and preferences. Evidence for the benefit of a particular intervention within a specific clinical context is identified through systematic review and is then balanced against the evidence for harms. In this regard, harms are broadly defined and may include serious adverse events, less serious adverse events that affect tolerability, minor adverse events, negative effects of the intervention on quality of life, barriers and inconveniences associated with treatment, direct and indirect costs of the intervention (including opportunity costs), and other negative aspects of the treatment that may influence decision making by the patient, the clinician, or both.

Many topics covered in this guideline have relied on forms of evidence such as consensus opinions of experienced clinicians or indirect findings from observational studies rather than research from randomized trials. It is well recognized that there are guideline topics and clinical circumstances for which high-quality evidence from clinical trials is not possible or is unethical to obtain (Council of Medical Specialty Societies 2012). For example, many questions need to be asked as part of an assessment and inquiring about a particular symptom or element of the history cannot be separated out for study as a discrete intervention. It would also be impossible to separate changes in outcomes due to assessment from changes in outcomes due to ensuing treatment. Research on psychiatric assessments and some psychiatric interventions can also be complicated by multiple confounding factors such as the

interaction between the clinician and the patient or the patient's unique circumstances and experiences. The GRADE working group and guidelines developed by other professional organizations have noted that a strong recommendation or "good practice statement" may be appropriate even in the absence of research evidence when sensible alternatives do not exist (Andrews et al. 2013; Brito et al. 2013; Djulbegovic et al. 2009; Hazlehurst et al. 2013). For each guideline statement, we have described the type and strength of the available evidence as well as the factors, including patient preferences, that were used in determining the balance of benefits and harms.

GUIDELINE SCOPE

The scope of this document is shaped by the diagnostic criteria for eating disorders and by the available evidence as obtained by a systematic review of the literature through September 2021, particularly focusing on AN, BN, and BED as defined by DSM-III, DSM-III-R, DSM-IV, DSM-IV-TR, DSM-5, or ICD-10. This practice guideline addresses evidence-based pharmacological, psychotherapeutic, and other nonpharmacological treatments for eating disorders in adolescents, emerging adults, and adults. In addition, it includes statements related to assessment and treatment planning, which are an integral part of patient-centered care.

Our systematic review attempted to include literature on avoidant/restrictive food intake disorder (ARFID); however, rigorous clinical trial data were not available due to the relative recency of the introduction of this diagnosis. We have included some discussion of ARFID in the implementation sections of this document, particularly as it relates to assessment and treatment planning. We specifically excluded rumination disorder and pica from our search of the literature due to their typical age of onset in infancy or childhood and the limited evidence on their treatment. We also excluded treatment of obesity from the scope of this guideline because obesity is not categorized as an eating disorder.

Data are also limited on individuals with eating disorders and significant physical health conditions or co-occurring psychiatric conditions, including substance use disorders. Many of the available studies of eating disorders did not analyze data separately for these patient subgroups or excluded individuals with these comorbidities. Nevertheless, in the absence of more robust evidence, the statements in this guideline should generally be applicable to individuals with co-occurring conditions. Additionally, although treatment-related costs are often barriers to receiving treatment and cost-effectiveness considerations are relevant to health care policy, cost-effectiveness considerations are outside the scope of this guideline.

The full text of the practice guideline describes aspects of guideline implementation that are relevant to individual patients' circumstances and preferences. A detailed description of research evidence related to the effects of pharmacological and nonpharmacological treatments in individuals with eating

disorders can be found in the appendices accompanying the full guideline at psychiatryonline.org/guidelines.

AUTHOR AND ARTICLE INFORMATION

Send correspondence to Jennifer Medicus (jmedicus@psych.org).

From the APA Practice Guideline Writing Group (Catherine Crone, M.D., Chair).

APA and the Guideline Writing Group especially thank Laura J. Fochtmann, M.D., M.B.I., Blair Uniacke, M.D., Seung-Hee Hong, and Jennifer Medicus for their outstanding work and effort in developing this guideline. APA also wishes to acknowledge the contributions of other APA staff including Michelle Dirst, Andrew Lyzenga, and Kristin Kroeger Ptakowski. APA wishes to give special recognition to Joel Yager, M.D. for his decades of contributions to APA and its practice guidelines, including his work on the Systematic Review Group and serving as Chair of the Eating Disorders Writing Group for three prior versions of this guideline. APA also thanks the APA Committee on Practice Guidelines (Daniel J. Anzia, M.D., Chair), liaisons from the APA Assembly for their input and assistance, and APA Councils and others for providing feedback during the comment period.

Practice Guidelines are assessments of scientific and clinical information that are current as of the date of authorship but are not continually updated and may not reflect the most recent evidence. They are provided as an educational service and should not be considered as a statement of the standard of care or inclusive of all proper treatments or methods of care. They are not intended to substitute for the independent professional judgment of the treating clinician. The ultimate recommendation regarding a particular assessment, clinical procedure, or treatment plan must be made by the clinician in light of the psychiatric evaluation, other clinical data, and the diagnostic and treatment options available. The guidelines are available on an "as is" basis, and APA makes no warranty, expressed or implied, regarding them. APA assumes no responsibility for any injury or damage to persons or property arising out of or related to any use of the guidelines.

Am J Psychiatry 2023; 180:167–171; doi: 10.1176/appi.ajp.23180001

REFERENCES

Agency for Healthcare Research and Quality: Methods guide for effectiveness and comparative effectiveness reviews. AHRQ Publ No 10(14)-EHC063-EF. Rockville, MD, Agency for Healthcare Research and Quality, 2014

Agostino H, Burstein B, Moubayed D, et al: Trends in the incidence of new-onset anorexia nervosa and atypical anorexia nervosa among youth during the COVID-19 pandemic in Canada. *JAMA Netw Open* 2021; 4:e2137395

Ahn J, Lee JH, Jung YC: Predictors of suicide attempts in individuals with eating disorders. *Suicide Life Threat Behav* 2019; 49:789–797

American Psychiatric Association: Practice Guideline for the Treatment of Patients with Eating Disorders, 3rd Ed. Arlington, VA, American Psychiatric Publishing, 2006

American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, 5th Ed, Text Revision. Washington, DC, American Psychiatric Association, 2022

Andrews JC, Schünemann HJ, Oxman AD, et al: GRADE guidelines: 15. Going from evidence to recommendation—determinants of a recommendation’s direction and strength. *J Clin Epidemiol* 2013; 66:726–735

Asch DA, Buresh J, Allison KC, et al: Trends in US patients receiving care for eating disorders and other common

behavioral health conditions before and during the COVID-19 pandemic. *JAMA Netw Open* 2021; 4:e2134913

Auger N, Potter BJ, Ukah UV, et al: Anorexia nervosa and the long-term risk of mortality in women. *World Psychiatry* 2021; 20:448–449

Balshem H, Helfand M, Schünemann HJ, et al: GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011; 64:401–406

Brito JP, Domecq JP, Murad MH, et al: The Endocrine Society guidelines: when the confidence cart goes before the evidence horse. *J Clin Endocrinol Metab* 2013; 98:3246–3252

Cliffe C, Shetty H, Himmerich H, et al: Suicide attempts requiring hospitalization in patients with eating disorders: a retrospective cohort study. *Int J Eat Disord* 2020; 53:458–465

Council of Medical Specialty Societies: Principles for the Development of Specialty Society Clinical Guidelines. Chicago, IL, Council of Medical Specialty Societies, 2012

Djulfbegovic B, Trikalinos TA, Roback J, et al: Impact of quality of evidence on the strength of recommendations: an empirical study. *BMC Health Serv Res* 2009; 9:120:120

Galmiche M, Déchelotte P, Lambert G, et al: Prevalence of eating disorders over the 2000–2018 period: a systematic literature review. *Am J Clin Nutr* 2019; 109:1402–1413

Gibbins NK, Kurdyak PA, Colton PA, et al: Diabetic ketoacidosis and mortality in people with type 1 diabetes and eating disorders. *Diabetes Care* 2021; 44:1783–1787

Guyatt G, Gutterman D, Baumann MH, et al: Grading strength of recommendations and quality of evidence in clinical guidelines: report from an American College of Chest Physicians Task Force. *Chest* 2006; 129:174–181

Guyatt G, Eikelboom JW, Akl EA, et al: A guide to GRADE guidelines for the readers of JTH. *J Thromb Haemost* 2013; 11:1603–1608

Hazlehurst JM, Armstrong MJ, Sherlock M, et al: A comparative quality assessment of evidence-based clinical guidelines in endocrinology. *Clin Endocrinol (Oxf)* 2013; 78:183–190

Institute of Medicine: Clinical Practice Guidelines We Can Trust. Washington, DC, National Academies Press, 2011

Kazdin AE, Fitzsimmons-Craft EE, Wilfley DE: Addressing critical gaps in the treatment of eating disorders. *Int J Eat Disord* 2017; 50:170–189

Keski-Rahkonen A: Epidemiology of binge eating disorder: prevalence, course, comorbidity, and risk factors. *Curr Opin Psychiatry* 2021; 34:525–531

Nielsen S, Vilmar JW: What can we learn about eating disorder mortality from eating disorder diagnoses at initial assessment? A Danish nationwide register follow-up study using record linkage, encompassing 45 years (1970–2014). *Psychiatry Res* 2021; 303:114091

Otto AK, Jary JM, Sturza J, et al: Medical admissions among adolescents with eating disorders during the COVID-19 pandemic. *Pediatrics* 2021; 148:e2021052201

Santomauro DF, Melén S, Mitchison D, et al: The hidden burden of eating disorders: an extension of estimates from the Global Burden of Disease Study 2019. *Lancet Psychiatry* 2021; 8:320–328

- Smith AR, Zuromski KL, Dodd DR: Eating disorders and suicidality: what we know, what we don't know, and suggestions for future research. *Curr Opin Psychol* 2018; 22:63–67
- Streatfeild J, Hickson J, Austin SB, et al: Social and economic cost of eating disorders in the United States: evidence to inform policy action. *Int J Eat Disord* 2021; 54:851–868
- Taquet M, Geddes JR, Luciano S, et al: Incidence and outcomes of eating disorders during the COVID-19 pandemic. *Br J Psychiatry* 2021; 220:1–3
- Tith RM, Paradis G, Potter BJ, et al: Association of bulimia nervosa with long-term risk of cardiovascular disease and mortality among women. *JAMA Psychiatry* 2020; 77:44–51
- Udo T, Grilo CM: Prevalence and correlates of DSM-5-defined eating disorders in a nationally representative sample of U.S. adults. *Biol Psychiatry* 2018; 84:345–354
- Udo T, Grilo CM: Psychiatric and medical correlates of DSM-5 eating disorders in a nationally representative sample of adults in the United States. *Int J Eat Disord* 2019; 52: 42–50
- Udo T, Bitley S, Grilo CM: Suicide attempts in US adults with lifetime DSM-5 eating disorders. *BMC Med* 2019; 17: 120
- van Hoeken D, Hoek HW: Review of the burden of eating disorders: mortality, disability, costs, quality of life, and family burden. *Curr Opin Psychiatry* 2020; 33: 521–527
- Wu J, Liu J, Li S, et al: Trends in the prevalence and disability-adjusted life years of eating disorders from 1990 to 2017: results from the Global Burden of Disease Study 2017. *Epidemiol Psychiatr Sci* 2020; 29:e191